

Ver. 0.94

Cat. No. **521 - 200** (reaction vol. 20 μl)
 521 - 500 (reaction vol. 50 μl)

Storage at : -20 °C

Lot. No. **A0000BCD**
Expiration date : 2000.00.00

Components

Taq PCR premix 96 tubes (0.2 ml 8-tube strip x 12ea)
 0.2 ml PCR tube storage rack

Description

Taq premix is made from geneall Taq DNA polymerase which is purified from the cloned *Thermos aquaticus* DNA polymerase gene expressed in *E. coli*. This mix contains all reaction components required for PCR, such as reaction buffer, dNTP, gel loading dye, stabilizer and sediment in addition to Taq DNA polymerase. It is recommended for use in routine PCR (below 10 kb), TA cloning and primer extension. This mix is stable for 1 year at -20 °C, 2 weeks at room temperature. It is lyophilized blue pellet type, so it is not necessary thawing time. It is controlled various reaction volume according to experience purpose. Therefore, this kit serve time-saving, cost-effective and good product. Included loading dye migrates through 1.0% agarose gels run in 0.5X TBE at approximately the same rate as DNA 300bp in length.

Storage condition

stable for 1 year at -20 °C, 2 weeks at room temperature.

PCR master mix components

(reaction vol. 20 μl , 50 μl)

Taq DNA polymerase	2U, 4U
dNTP (mixture)	200 μM (800 μM)
reaction buffer (2.5 mM MgCl ₂)	1X
loading dye & stabilizer	1X

Features

high efficiency
 ready to use
 stable for 2 weeks at room temperature (1 year at -20 °C)
 stable to thawing at 5 times

Application

general PCR reaction
 TA-cloning

Reaction mixture (example)

reaction vol.	20 μl	50 μl
Taq premix	- μl	- μl
primer 1 (10 pmol/ μl)	1 μl	1-2 μl
primer 2 (10 pmol/ μl)	1 μl	1-2 μl
template	1-50 ng	1-100 ng
DW	up to 20 μl	up to 50 μl
final reaction vol.	20 μl	50 μl

Thermal cycling condition

95 °C	2 min	} 30 - 35 cycles
95 °C	20 sec	
A °C	10 sec	
72 °C	B min	
72 °C	2-5 min	

A : The value is 4~6 lower than T_m of primers

$$T_m = 2(A+T) + 4(G+C)$$

B : below 3 kb 0.5 - 1 min/kb
 more than 3 kb 1 - 2 min/kb

HQ buffer

- In GC-rich reaction, HQ buffer increases the activity of Taq DNA polymerase.
- HQ buffer removes a hair-pin structure of GC-rich region.
- The dilution factor of HQ buffer is variable, 0.5X - 2X, depending on a case by case basis.
- We recommend to use of HQ buffer in PCR reaction of long-size target.

HQ buffer (example)

reaction vol.	20 μl	50 μl
0.5X HQ	2 μl	5 μl
1X HQ	4 μl	10 μl
1.5X HQ	6 μl	15 μl
2X HQ	8 μl	20 μl

End note : For research use only. Not for use in diagnostic or therapeutic procedures.